

List of Essential Medicines for sixth-year students.  
Chair of Internal Disease №1 with a course of endocrinology and hematology

1. Glitserintrinitrate (nitroglycerin) (Table, caps., For injection)

Rp.: Nitroglycerini 0,0005

D. td № 50 in tab. (Caps.)

Table S. 1. (Kaps.) under the tongue at an acute attack of angina.

Rp.: Sol. Nitroglycerini 0,1%-100ml.

D. td 1

S. administered intravenously at 10 mg. per minute.

2. Isosorbidemononitrate (Table, caps.)

Rp.: Isosorbidimononitrati 0,04

D. td № 60 in tab. (Caps.)

Table S. 1. (Kaps.), 2 times a day.

3. Isosorbidedinitrate (Table, caps.)

Rp.: Isosorbididinitrati 0,04

D. td № 60 in tab. (Caps.)

Table S. 1. (Kaps.), 2 times a day.

4. Propranolol (Table)

Rp.: Propranololi 0,04

D. td № 60 in tab.

Table S. 1. 2 times a day.

5. Atenolol (Table)

Rp.: Atenololi 0,05

D. td № 60 in tab.

Table S. 1. 2 times a day.

6. Metoprolol (Table)

Rp.: Metoprololi 0,05

D. td № 60 in tab.

Table S. 1. 2 times a day.

7. Bisoprolol (Table)

Rp.: Bisoprololi 0,005

D. td № 30 in tab.

Table S. 1. 1 times a day.

8. Carvedilol (Table, caps.)

Rp.: Carvediloli 0,025

D. td № 60 in tab.

Table S. 1. (Kaps.) 2 times a day.

9. Verapamil (Table, caps., For injection)

Rp.:Verapamili 0,08 D. td № 60 in tab.

Table S. 1. (Kaps.)2 times a day.

Rp.: Sol. Verapamili 0,25% - 2ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

10. Amlodipine (Table)

Rp.:Amlodipini 0,01

D. td № 30 in tab.

Table S. 1. 1 times a day.

11. Nifedipine (Table)

Rp.:Nifedipini 0,01

D. td № 30 in tab.

Table S. 1. under the tongue at high pressure.

12. Procainamide (procainamide) (for injection)

Rp.: Sol. Novocainamidi 10% - 5ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

13. Lidocaine (for injection)

Rp.: Sol. Lidocaini 2% - 2ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

14. Amiodarone (Table, for injection)

Rp.:Amiodaroni 0,2

D. td № 30 in tab.

Table S. 1. 1 times a day.

Rp.: Sol. Amiodaroni 5% - 3ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

15. Sotalol (Table)

Rp.:Sotaloli 0,08

D. td № 60 in tab.

Table S. 1. 2 times a day.

16. Dopamine (for injection)

Rp.: Sol. Dopamini 4% - 5ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

17. Epinephrine (adrenaline) (for injection)  
Rp.: Sol. Adrenalinihydrochloridi 0,1% - 1ml.  
D. td № 1 in ampull.  
S. enter subcutaneously with 0.5 ml.

18. Strofantin (for injection)  
Rp.: Sol. Strophantini K 0,025% - 1ml.  
D. td № 1 in ampull.  
S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

19. Digoxin (Table for injection)  
Rp.:Digoxini 0,00025  
D. td № 50 in tab.  
Table S. 1. 1 times a day.  
Rp.: Sol. Digoxini 0,025%-1ml.  
D. td № 1 in ampull.  
S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

20. Captopril (Table)  
Rp.:Captoprili 0,05  
D. td № 60 in tab.  
Table S. 1. 3 times per day.

21. Enalapril (Table)  
Rp.:Enalaprili 0,01  
D. td № 60 in tab.  
Table S. 1. 2 times a day.

22. Lisinopril (Table)  
Rp.:Lizinoprili 0,01  
D. td № 60 in tab.  
Table S. 1. 2 times a day.

23. Fosinopril (Table)  
Rp.:Fozinoprili 0,01  
D. td № 60 in tab.  
Table S. 1. 2 times a day.

24. Ramipril (Table)  
Rp.:Ramiprili 0,005  
D. td № 60 in tab.  
Table S. 1. 2 times a day.

25. Losartan (Table)

Rp.:Losartani 0,05

D. td № 30 in tab.

Table S. 1. 1 times a day.

26. Trimetazidine (Table)

Rp.:Trimetazidini 0,035

D. td № 60 in tab.

Table S. 1. 2 times a day.

27. Furosemide (Table for injection)

Rp.:Furosemidi 0,04

D. td № 30 in tab.

Table S. 1. 1 time per day in the morning.

Rp.: Sol. Furosemidi 1% - 2ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

28. Indapamide (Table)

Rp.:Indapamidi 0,0025

D. td № 30 in tab.

Table S. 1. 1 in the morning every day.

29. Hydrochlorothiazide (Table)

Rp.:Hydrochlorthiazidi 0,025

D. td № 30 in tab.

Table S. 1. 1 time per day in the morning.

30. Spironalakton (veroshpiron) (Table, caps.)

Rp.:Spironolactoni 0,05

D. td № 30 in tab. (Caps.)

Table S. 1. (Kaps.)1 times a day, morning.

31. Clonidine (clonidine) (Table, for injection)

Rp.:Clophelini 0,00015

D. td № 30 in tab.

Table S. 1. under the tongue at high pressure.

Rp.: Sol. clophelini 0,01% - 1ml.

D. td № 1 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution slowly.

32. Morphine (for injection)

Rp.: Sol. Morphini 1% - 1ml.

D. td № 1 in ampull.

S. enter subcutaneously 1 ml.

33. Tramadol (Table, caps., For injection)

Rp.:Tramadol 0,05

D. td № 10 in tab. (Caps.) Table

S. 1. (Kaps.), 2 times a day. Rp.:

Sol. Tramadol 5% - 2ml. D. td

№ 1 in ampull.

1 S. Introducing intramuscularly once a day.

34. Heparin (for injection)

Rp.: Sol. Heparini5ml (1ml - 5000ME)

D. td № 1 in ampull.

S. enter subcutaneously 1 ml into the anterolateral area of the abdominal wall 4 times a day.

35. Enoxaparin (Clexane) (for injection)

Rp.: Sol. Klexani 10% - 1 ml (1ml - 10000ME)

D. td № 1

S. injected subcutaneously in the region of the anterolateral abdominal wall 4 times per day.

36. Warfarin (Table)

Rp.:Warfarini 0,0025

D. td № 50 in tab.

S. According to Table 2. 1 time per day in the evening.

37. Streptokinase (lyophilized powder for solution for injection, infusion)

Rp.:Streptokinasi 250000 ED

D. td № 1

S. contents of the vial to dissolve in 100ml. 0.9% sodium chloride solution administered intravenously.

38. Alteplase (Micardis) (lyophilized powder for solution for injection, infusion)

Rp.:Alteplasi 0,05

D. td № 1

S. contents of the vial to dissolve in 100ml. 0.9% sodium chloride solution administered intravenously.

39. Pentoxifylline (Trental) (Table, caps.)

Rp.:Pentoxiphyllini 0,2

D. td № 60 in tab. (Caps.)

Table S. 1. (Kaps.), 2 times a day.

40. Dipyridamole (chimes) (Table)

Rp.:Dipiridamoli 0,025

D. td № 90 in tab.

Table S. 1. 3 times per day.

41. Clopidogrel (Plavix) (Table)

Rp.:Clopidogreli 0,075

D. td № 28 in tab.

Table S. 1. 1 times a day.

42. Simvastatin (Table)

Rp.:Simvastatini 0,02

D. td № 30 in tab.

Table S. 1. one evening after dinner every day.

43. Lovastatin (Table)

Rp.:Lovastatini 0,02

D. td № 30 in tab.

Table S. 1. one evening after dinner every day.

44. Atorvastatin (Table)

Rp.:Atorvastatini 0,02

D. td № 30 in tab.

Table S. 1. one evening after dinner every day.

45. Indomethacin (Table ointment, gel)

Rp.:Indometacini 0,025

D. td № 30 in tab.

Table S. 1. 3 times per day.

Rp.: Ung. Indometacini 10% - 30.0

D. td № 1

S. topically to the affected joint area two times daily.

Rp.: Gel. Indometacini 1% - 50.0

D. td № 1

S. topically to the affected joint area two times daily.

46. Acetylsalicylic acid (aspirin) (Table)

Rp.:Ac.acetylsalicylici 0,5

D. td № 10 in tab.

Table S. 1. 3 times per day.

47. Diclofenac (sodium salt) (Table, caps., Gel for injection)

Rp.:Diclofenaci 0,025

D. td № 10 in tab.

Table S. 1. 2 times a day.

Rp.:Diclofenaci 0,1

D. td № 10 in caps.

S. 1 caps. 1 times a day.

Rp.: Sol. Diclofenaci 2,5% - 3ml. D.

td № 5 in ampull.

1 S. Introducing intramuscularly once a day.

Rp.: Gel. Diclofenaci 1% - 50.0

D. td № 1

S. topically to the affected joint area two times daily.

48. Ibuprofen (Table)

Rp.:Ibuprofeni 0,2

D. td № 20 in tab.

Table S. 1. 3 times per day.

49. Meloxicam (Table for injection)

Rp.:Meloxicami 0,0075

D. td № 10 in tab.

Table S. 1. 1 times a day.

Rp.: Sol. Meloxicami 0,015 - 1,5 ml. D.

td № 3 in ampull.

1 S. Introducing intramuscularly once a day.

50. Nimesulide (Table powders)

Rp.:Nimesulidi 0,1

D. td № 10 in tab. (Pulv.)

Table S. 1. (Powder) two times daily; Filter powder dissolved in 1/2 cup warm water.

51. Paracetamol (Table)

Rp.:Paracetamoli 0,5

D. td № 10 in tab.

Table S. 1. 3 times per day.

52. Hydroxychloroquine (Table)

Rp.:Hydroxychloroquini 0,2

D. td № 100 in tab.

Table S. 1. 2 times a day.

53. Methotrexate (Table)

Rp.:Methotrexati 0,0025

D. td № 20 in tab.

Table S. 1. 3 times per week.

54. Azathioprine (Table)

Rp.:Azathioprini 0,05

D. td № 25 in tab.

Table S. 1. 2 times a day.

55. Cyclophosphamide (Table, for injection)

Rp.:Cyclophosphamidi 0,05

D. td № 10 in tab.

Table S. 1. 1 times a day.

Rp.:Cyclophosphamidi 0,1

D. td № 1

S. contents of the vial to dissolve in water for injection administered intramuscularly 3 times a day.

56. Leflunomide (Table)

Rp.:Leflunomidi 0,02

D. td № 30 in tab.

Table S. 1. 1 times a day.

57. Sulfasalazine (Table)

Rp.:Sulfasalazini 0,5

D. td № 50 in tab.

Table S. 1. 4 times a day.

58. Penicillamine (Table)

Rp.:Penicillamini 0,25

D. td № 100 in tab.

Table S. 1. 1 times a day.

59. Allopurinol (Table)

Rp.:Allopurinoli 0,1

D. td № 30 in tab.

Table S. 1. 1 times a day.

60. Chondroitin sulfate (kaps. for injection)

Rp.:Chondroitinisulfati 0,25

D. td № 90 in caps.

S. 1 caps. 3 times per day.

Rp.: Sol. Chondroitinisulfati 0,1 - 1ml.

D. td № 20 in ampull.

S. Introducing 2ml intramuscularly. every other day.

61. Alendronate (Table)

Rp.:Ac.alendronici 0,07

D. td № 4 in tab.

Table S. 1. 1 time per week.

62. Prednisolone (Table for injection)

Rp.:Prednisoloni 0,005

D. td № 30 in tab.

Table S. 1. 1 times a day.

Rp.:Prednisolonihemisuccinati 0,025

D. td № 1

S. dissolve the contents of the vial with 20 ml. 0.9% sodium chloride solution administered intravenously.

63. Methylprednisolone (Table for injection)

Rp.:Methylprednisoloni 0,004

D. td № 30 in tab.

Table S. 1. 1 times a day.

Rp.:Susp. methylprednisoloni 4% - 2ml.

D. td № 5 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

64. Dexamethasone (Table for injection)

Rp.:Dexamethasoni 0,0005

D. td № 30 in tab.

Table S. 1. 1 times a day.

Rp.: Sol. dexamethasoni 0,4% - 2ml.

D. td № 5 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

65. Atropine Sulfate (for injection)

Rp.: Sol. Atropinisulfati 0,1% - 1ml.

D. td № 5 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

66. Aminophylline (Table for injection)

Rp.:Aminophyllini 0,15

D. td № 30 in tab.

Table S. 1. 3 raza per day.

Rp.: Sol. Aminophyllini 2,4% - 5ml.

D. td № 5 in ampull.

S. Introducing intravenously with 20 ml. 0.9% sodium chloride solution.

67. Drotaverinum (Nospanum) (Table, for injection)

Rp.:Drotaverini 0, 04

D. td № 30 in tab.

Table S. 1. 3 raza per day.

Rp.: Sol. Drotaverini 2% - 2ml.

D. td № 5 in ampull.

S. Introducing intramuscularly 2 times a day.

68. Papaverine (for injection)

Rp.: Sol. Papaverini 2% - 2ml.

D. td № 5 in ampull.

S. Introducing intramuscularly 2 times a day.

69. Metoklopromid (Table for injection)

Rp.:Metoclopromidi 0,01

D. td № 30 in tab.

Table S. 1. 3 raza per day.

Rp.: Sol. metoclopromidi 0,5% - 2ml.

D. td № 5 in ampull.

S. Introducing intramuscularly 3 times a day.

70. Multienzymy comprising amylase, lipase, protease (pancreatin)

(Table)

Rp.: Tab. «Pancreatin» № 90

DS 1 tab. 3 raza a day 20 minutes before meals.

71. Mesalazine (Table) Rp.:Mesalazini 0,5

D. td № 50 in tab.

Table S. 1. 3 raza per day.

72. Ursodeoxycholic acid (ursosan, ursokaps) (kaps.)

Rp.: Ac. ursodeoxycholici 0 25

D. td № 50 in caps.

S. 2 caps. 2 at bedtime, without chewing.

73. Lactulose (Duphalac) (syrup)

Rp.: Sir. Lactulosi 50% - 200ml.

D. td № 1

S. 1 tablespoon 3 times a day.

74. Ademetionine (Table, for injection)

Rp.:Ademethionini 0,4

D. td № 20 in tab.

Table S. 1. 2 times a day.

Rp.:Ademethionini 0,4

D. td № 1

S. contents of the vial to dissolve in 250 ml. 0.9% sodium chloride solution administered intravenously.

75. Essential phospholipids (essentiale H) (kaps.)

Rp.: Caps. «Essentiale H» »№ 50 DS

1 caps. 3 raza per day.

76. Omeprazole (kaps.)

Rp.:Omeprasoli 0, 02

D. td № 60 in caps.

S. 1 caps. 2 times a day.

77. Ranitidine (Table for injection)

Rp.:Ranitidini 0, 15

D. td № 60 in tab.

Table S. 1. 2 times a day.

Rp.: Sol. ranitidini 1% - 5ml.

D. td № 5 in ampull.

S. Introducing intramuscularly 2 times a day.

78.Famotidine (Table)

Rp.:Famotidini 0,04

D. td № 30 in tab.

Table S. 1. 2 times a day.

79. Loperamide (kaps.)

Rp.:Loperamidi 0, 02

D. td № 20 in caps.

S. 1 caps. 3 raza per day.

80. Folic acid (Table)

Rp.: Ac. folici 0, 001

D. td № 50 in tab.

Table S. 1. 3 raza per day.

81. Ferrous sulfate (tardiferon, ferrogradument) (Table)

Rp.:Ferrosisulfati 0, 325

D. td № 90 in tab.

Table S. 1. 3 raza per day.

82. Tocopherol acetate (kaps.)

Rp.:Tocopheroliacetati 0, 1

D. td № 90 in caps.

S. 1 caps. 3 raza per day.

83. Cyanocobalamin (for injection)

Rp.: Sol. Cyanocobalamin 0,1% - 1ml.

D. td № 5 in ampull.

1 S. Introducing intramuscularly once a day.

84. Etamsylate (Table for injection)

Rp.:Etamsylati 0, 05

D. td № 100 in tab.

Table S. 1. 3 raza per day.

Rp.: Sol. etamsylati 12,5% - 2ml.

D. td № 5 in ampull.

S. injected intravenously with 200 ml. 0.9% sodium chloride solution.

85. Aminocaproic acid (for injection)  
Rp.: Sol. Ac. aminocapronici 5% - 100ml.  
D. td № 1  
S. administered intravenously.

86. Amoxicillin (kaps., table).  
Rp.:Amoxicillini 0 5  
D. td № 20 in tab. (Caps.) Table  
S. 1. (Kaps.)3 raza day.

87. Ampicillin (Table for injection)  
Rp.:Ampicillini 0 5  
D. td № 20 in tab.  
Table S. 1. 4 times a day.  
Rp.:Ampicillini 0,5  
D. td № 20  
S. contents of the vial diluted in 3 ml. water for injection, intramuscular injection four times per day.

88. Amoxicillin / clavulanic acid (for injection, infusion, Table).  
Rp.:Amoxicillinclavulonati 0, 625  
D. td № 20 in tab.  
Table S. 1. 3 raza per day.  
Rp.:Amoxicillinclavulonati 1,2  
D. td № 10  
S. contents of the vial to dissolve in 20 ml. 0.9 sodium chloride solution, intravenously administered two times a day.

89. Ceftriaxone (for injection)  
Rp.:Ceftriaxoni 1,0  
D. td № 20  
S. contents of the vial diluted in 3 ml. 0, 5% novocaine administered intramuscularly, 2 times a day.

90. Cefotaxime (for injection)  
Rp.:Cefotaximi 1,0  
D. td № 20  
S. contents of the vial diluted in 3 ml. 0, 5% novocaine administered intramuscularly, 2 times a day.

91. Cefepime (for injection)  
Rp.:Cefepimi 1,0  
D. td № 20  
S. contents of the vial diluted in 3 ml. 0, 5% novocaine administered intramuscularly, 2 times a day.

92. Amikacin (for injection)  
Rp.:Amikacinisulfati 0,5

D. td № 20

S. contents of the vial diluted in 3 ml. 0, 5% novocaine administered intramuscularly 3 times a day.

93. Ciprofloxacin (Table, caps., For injection)

Rp.:Ciprofloxacini 0 5

D. td № 20 in tab. (Caps.)

Table S. 1. (Kaps.)2 times a day.

Rp.: Sol. ciprofloxacini 0,2% - 100ml.

D. td № 1

S. 1 administered intravenously once a day.

94. Ofloxacin (infusion, Table., Caps.)

Rp.:Ofloxacini 0 2

D. td № 10 in tab. (Caps.) Table

S. 1. (Kaps.)2 times a day.

Rp.: Sol. ofloxacini 0,2% - 100ml.

D. td № 1

S. 1 administered intravenously once a day.

95. Levofloxacin (Table, caps.)

Rp.:Levofloxacini 0 5

D. td № 10 in tab. (Caps.)

Table S. 1. (Kaps.)2 times per day.]

96. Meropenem (for injection)

Rp.:Meropenemi 1,0

D. td № 20

S. contents of the vial diluted in 3 ml. 0, 5% novocaine administered intramuscularly 3 times a day.

97. Nitroxoline (Table)

Rp.:Nitroxolini 0, 05

D. td № 50 in tab.

S. According to Table 2. 4 times a day.

98. Metronidazole (for infusion, Table.,)

Rp.:Metronidazoli 0, 25

D. td № 20 in tab.

By S. 1tabl. 4 times a day.

Rp.:Metronidazolihemisuccinati 0,5

D. td № 20

S. contents of the vial to dissolve in 250 ml. 0.9% sodium chloride solution, administered intravenously once a day 1.

99. Azithromycin (Table, caps.)

Rp.:Azithromycini 0 5  
D. td № 3 in tab. (Caps.)  
Table S. 1. (Kaps.) 1 times a day.

100. Clarithromycin (Table)  
Rp.:Clarithromycini 0 5  
D. td № 7 in tab.  
Table S. 1. 1 times a day.