

1. Hypothyroidism is more common for:
  - a) male
  - b) women
  - c) children
  - d) any people
  
2. In primary hypothyroidism the TSH will be:
  - a) increased
  - b) decreased
  - c) does not change
  
3. The average replacement dose of levothyroxine sodium for patients with hypothyroidism is:
  - a) 1.0 - 1.2  $\mu\text{g}$  / kg / day
  - b) 1.6 - 1.8  $\mu\text{g}$  / kg / day
  - a) 2.0 - 2.2  $\mu\text{g}$  / kg / day
  
4. In the appointment of replacement therapy in women is not recommended:
  - a) pregnant for at least 1 year
  - b) to breastfeed
  - c) there are no restrictions
  
5. The main reason for hyperthyroiditis in young adults:
  - a) excessive intake of iodine in the body
  - b) Graves disease
  - c) thyroid cancer
  
6. In primary thyrotoxicosis TSH will be:
  - a) reduced
  - b) increased
  - c) has not changed
  
7. The main method of diagnosis of Graves' disease is:
  - a) determination of TSH and free T<sub>4</sub>+TSH<sub>r</sub>-Ab
  - b) thyroid biopsy
  - c) thyroid scintigraphy
  
8. Indications for surgical treatment of diffuse toxic goiter:
  - a) all patients with Graves disease
  - b) the ineffectiveness of prolonged conservative treatment
  - c) the development of side effects with conservative treatment
9. The major complications of antithyroid therapy is:
  - a) agranulocytosis
  - b) cutaneous manifestations
  - c) thyroid cancer
  
10. Diagnostic criteria for subacute thyroiditis:
  - a) pain in the thyroid gland
  - b) a significant increase in ESR
  - c) sub-febrile fever